

## CLAIMS

We claim:

1. A composition (for inhibition of inducible COX-2) activity and having minimal effect on COX-1 activity, said composition comprising, as a first component an effective amount of a sesquiterpene lactone species and an effective amount of a second component selected from the group consisting of a diterpene lactone species and a triterpene species or derivatives thereof.

2. The composition of Claim 1 wherein first and second components are derived from plants or plant extracts.

3. The composition of Claim 1 wherein at least one of said first or second component is conjugated with a compound selected from the group consisting of mono- or disaccharides, amino acids, sulfates, succinate, acetate and glutathione.

4. The composition of Claim 1, formulated in a pharmaceutically acceptable carrier.

5. The composition of Claim 1, additionally containing one or members selected from the group consisting of antioxidants, vitamins, minerals, proteins, fats, carbohydrates, glucosamine, chondroitin sulfate and aminosugars.

5 603  
6. A composition for inhibition of inducible COX-2 activity and having minimal effect on COX-1 activity, said composition comprising, as a first component an effective amount of a pharmaceutical grade compound selected from the group consisting of parthenolide, encelin, leucanthin B, enhydrin, melapodin A, tenulin, confertiflorin, burrodin, psilostachyin A, costunolide, strigol and <sup>terpene</sup> helenalin; and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, dehydroandrographolide, deoxyandrographolide, aneoandrographolide, ursolic acid, oleanolic acid, betulin, betulinic acid, glycyrrhetic acid, glycyrrhizic acid, triperin and derivatives thereof.

10

7. The composition of Claim 6 wherein first and second components are derived from plants or plant extracts.

15 603  
8. The composition of Claim 6 wherein at least one of said first or second component is conjugated with a compound selected from the group consisting of mono- or di- saccharides, amino acids, sulfates, succinate, acetate and glutathione.

9. The composition of Claim 6, formulated in a pharmaceutically acceptable carrier.

10. The composition of Claim 6, additionally containing one or members selected from the group consisting of antioxidants, vitamins, minerals, proteins, fats, carbohydrates, glucosamine, chondroitin sulfate and aminosugars.

11. A composition for inhibition of inducible COX-2 activity and having minimal effect on COX-1 activity, said composition comprising, as a first component an effective amount of a pharmaceutical grade compound selected from the group consisting of parthenolide, encelin, leucanthin B, enhydrin, and melapodin A; and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, dehydroandrographolide, deoxyandrographolide, neoandrographolide, ursolic acid, oleanolic acid, betulin, betulinic acid, glycyrrhetic acid, glycyrrhizic acid, triperin and derivatives thereof.

12. The composition of Claim 11 wherein first and second components are derived from plants or plant extracts.

13. The composition of Claim 11 wherein at least one of said first or second component is conjugated with a compound selected from the group consisting of mono- or di- saccharides, amino acids, sulfates, succinate, acetate and glutathione.

14. The composition of Claim 11, formulated in a pharmaceutically acceptable carrier.

15. The composition of Claim 11, additionally containing one or members selected from the group consisting of antioxidants, vitamins, minerals, proteins, fats, carbohydrates, glucosamine, chondroitin sulfate and aminosugars.

16. A composition for inhibition of inducible COX-2 activity and having minimal effect on COX-1 activity, said composition comprising, as a first component an effective amount of a pharmaceutical grade parthenolide and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, ursolic acid, oleanolic acid, and derivatives thereof.

17. The composition of Claim 16 wherein first and second components are derived from plants or plant extracts.

18. The composition of Claim 16 wherein at least one of said first or second component is conjugated with a compound selected from the group consisting of mono- or di- saccharides, amino acids, sulfates, succinate, acetate and glutathione.

20

19. The composition of Claim 16, formulated in a pharmaceutically acceptable carrier.

20. The composition of Claim 16, additionally containing one or members selected from the group consisting of antioxidants, vitamins, minerals, proteins, fats, carbohydrates, glucosamine, chondroitin sulfate and aminosugars.

21. A method of dietary supplementation in animals comprising administering to an animal suffering symptoms of inflammation a composition comprising, as a first component an effective amount of a sesquiterpene lactone species and an effective amount of a second component selected from the group consisting of a diterpene lactone species and a triterpene species or derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

22. The method of Claim 21 wherein the composition is formulated in a dosage form such that said administration provides from 0.05 to 5.0 mg body weight per day of each sequesterpene lactone species, and from 0.5 to 20.0 mg/kg bodyweight per day of each diterpene lactone species or triterpene species.

23. The method of Claim 21, wherein the composition is administered in an amount sufficient to maintain a serum concentration of 0.001 to 10  $\mu$ M of each sesquiterpene lactone species and from 0.001 to 10  $\mu$ M of each diterpene lactone or triterpene species.

24. The method of Claim 21 wherein said animal is selected from the group consisting of humans, non-human primates, dogs, cats, birds, horses and ruminants.

25. The method of Claim 21 wherein administration is by a means selected from the group consisting of oral, parenteral, topical, transdermal and transmucosal delivery.

26. A method of dietary supplementation in animals comprising administering to an animal suffering symptoms of inflammation a composition comprising, as a first component an effective amount of a pharmaceutical grade compound selected from the group consisting of parthenolide, encelin, leucanthin B, enhydrin, melapodin A, tenulin, confertiflorin, burrodin, psilostachyin A, costunolide, strigol and helenalin; and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, dehydroandrographolide, deoxyandrographolide, aneoandrographolide, ursolic acid, oleanolic acid, betulin, betulinic acid, glycyrrhetic acid, glycyrrhizic acid, triperin and derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

27. The method of Claim 26 wherein the composition is formulated in a dosage form such that said administration provides from 0.05 to 5.0 mg body weight per day of each sequesterpene lactone species, and from 0.5 to 20.0 mg/kg bodyweight per day of each diterpene lactone species or triterpene species.

5

28. The method of Claim 26, wherein the composition is administered in an amount sufficient to maintain a serum concentration of 0.001 to 10  $\mu$ M of each sesquiterpene lactone species and from 0.001 to 10  $\mu$ M of each diterpene lactone or triterpene species.

29. A method of dietary supplementation in animals comprising administering to an animal suffering symptoms of inflammation a composition comprising, as a first component an effective amount of a pharmaceutical grade compound selected from the group consisting of parthenolide, encelin, leucanthin B, enhydrin, and melapodin A; and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, dehydroandrographolide, deoxyandrographolide, neoandrographolide, ursolic acid, oleanolic acid, betulin, betulinic acid, glycyrrhetic acid, glycyrrhizic acid, triperin and derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

30. A method of dietary supplementation in animals comprising administering to an animal suffering symptoms of inflammation a composition comprising, as a first

component an effective amount of a pharmaceutical grade parthenolide and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, ursolic acid, oleanolic acid, and derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

31. A method of therapeutic treatment in animals comprising administering to an animal suffering symptoms of arthritis a composition comprising, as a first component an effective amount of a sesquiterpene lactone species and an effective amount of a second component selected from the group consisting of a diterpene lactone species and a triterpene species or derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

32. A method of therapeutic treatment in animals comprising administering to an animal suffering symptoms of arthritis a composition comprising, as a first component an effective amount of a pharmaceutical grade compound selected from the group consisting of parthenolide, encelin, leucanthin B, enhydrin, melapodin A, tenulin, confertiflorin, burrodin, psilostachyin A, costunolide, strigol and helenalin; and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, dehydroandrographolide, deoxyandrographolide, neoandrographolide, ursolic acid, oleanolic acid, betulin, betulinic acid, glycyrrhetic acid, glycyrrhizic acid,



triperin and derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

33. A method of therapeutic treatment in animals comprising administering to an animal suffering symptoms of arthritis a composition comprising, as a first component an effective amount of a pharmaceutical grade compound selected from the group consisting of parthenolide, encelin, leucanthin B, enhydrin, and melapodin A; and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, dehydroandrographolide, deoxyandrographolide, neoandrographolide, ursolic acid, oleanolic acid, betulin, betulinic acid, glycyrrhetic acid, glycyrrhizic acid, triperin and derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

34. A method of therapeutic treatment in animals comprising administering to an animal suffering symptoms of arthritis a composition comprising, as a first component an effective amount of a pharmaceutical grade parthenolide and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, ursolic acid, oleanolic acid, and derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

35. A method of therapeutic treatment comprising applying to the skin of a human suffering symptoms of acne rosacea a lotion comprising a composition comprising, as a first component an effective amount of a sesquiterpene lactone species and an effective amount of a second component selected from the group consisting of a diterpene lactone species and a triterpene species or derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

36. A method of therapeutic treatment comprising applying to the skin of a human suffering symptoms of acne rosacea a lotion comprising a composition comprising, as a first component an effective amount of a pharmaceutical grade compound selected from the group consisting of parthenolide, encelin, leucanthin B, enhydrin, melapodin A, tenulin, confertiflorin, burrodin, psilostachyin A, costunolide, strigol and helenalin; and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, dehydroandrographolide, deoxyandrographolide, neoandrographolide, ursolic acid, oleanolic acid, betulin, betulinic acid, glycyrrhetic acid, glycyrrhizic acid, triperin and derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

37. A method of therapeutic treatment comprising applying to the skin of a human suffering symptoms of acne rosacea a lotion comprising a composition comprising, as a first component an effective amount of a pharmaceutical grade compound selected from the

group consisting of parthenolide, encelin, leucanthin B, enhydrin, and melapodin A; and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, dehydroandrographolide, deoxyandrographolide, neoandrographolide, ursolic acid, oleanolic acid, betulin, betulinic acid, glycyrrhetic acid, glycyrrhizic acid, triperin and derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

5

38. A method of therapeutic treatment comprising applying to the skin of a human suffering symptoms of acne rosacea a lotion comprising a composition comprising, as a first component an effective amount of a pharmaceutical grade parthenolide and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, ursolic acid, oleanolic acid, and derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

10

39. A method of therapeutic treatment comprising applying to the skin of a human suffering symptoms of psoriasis a lotion comprising a composition comprising, as a first component an effective amount of a sesquiterpene lactone species and an effective amount of a second component selected from the group consisting of a diterpene lactone species and a triterpene species or derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

20

5  
10  
20  
30  
40  
50  
60  
70  
80  
90  
100  
110  
120  
130  
140  
150  
160  
170  
180  
190  
200  
210  
220  
230  
240  
250  
260  
270  
280  
290  
300  
310  
320  
330  
340  
350  
360  
370  
380  
390  
400  
410  
420  
430  
440  
450  
460  
470  
480  
490  
500  
510  
520  
530  
540  
550  
560  
570  
580  
590  
600  
610  
620  
630  
640  
650  
660  
670  
680  
690  
700  
710  
720  
730  
740  
750  
760  
770  
780  
790  
800  
810  
820  
830  
840  
850  
860  
870  
880  
890  
900  
910  
920  
930  
940  
950  
960  
970  
980  
990  
1000

40. A method of therapeutic treatment comprising applying to the skin of a human suffering symptoms of psoriasis a lotion comprising a composition comprising, as a first component an effective amount of a pharmaceutical grade compound selected from the group consisting of parthenolide, encelin, leucanthin B, enhydrin, melapodin A, tenulin, confertiflorin, burrocin, psilostachyin A, costunolide, strigol and helenalin; and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, dehydroandrographolide, deoxyandrographolide, neoandrographolide, ursolic acid, oleanolic acid, betulin, betulinic acid, glycyrrhetic acid, glycyrrhizic acid, triperin and derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

41. A method of therapeutic treatment comprising applying to the skin of a human suffering symptoms of psoriasis a lotion comprising a composition comprising, as a first component an effective amount of a pharmaceutical grade compound selected from the group consisting of parthenolide, encelin, leucanthin B, enhydrin, and melapodin A; and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, dehydroandrographolide, deoxyandrographolide, neoandrographolide, ursolic acid, oleanolic acid, betulin, betulinic acid, glycyrrhetic acid, glycyrrhizic acid, triperin and derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

42. A method of therapeutic treatment comprising applying to the skin of a human suffering symptoms of psoriasis a lotion comprising a composition comprising, as a first component an effective amount of a pharmaceutical grade parthenolide and a second component an effective amount of a pharmaceutical grade compound selected from the group consisting of andrographolide, ursolic acid, oleanolic acid, and derivatives thereof, and continuing said administering of the composition until said symptoms are reduced.

5

091506.073104